

Ramik[®] Green

(EPA Reg. No. 61282-46)

RODENTICIDE FOR CONTROL OF GROUND SQUIRRELS USING BAIT STATIONS

SPECIAL LOCAL NEED REGISTRATION FOR USE ONLY IN THE STATE OF MONTANA

ACTIVE INGREDIENT:

Diphacinone

(2-Diphenylacetyl-1,3-Indandione).....0.005%

INERT INGREDIENTS:.....99.995%

TOTAL:.....100.000%

EPA SLN No. MT-10-0002

Reg. No. 61282-46

EPA Est. No. 61282-WI-01

CAUTION

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION – Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

ENVIRONMENTAL HAZARDS – This product is toxic to mammals and birds. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Notice: It is a federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. The use of this product may pose a hazard to Federally designated endangered/ threatened species. Suggested measures to protect endangered species in your county may exist in an EPA “Interim Measures” pamphlet. You may call EPA’s Endangered Species Hotline, 1-800-447-3813, to find out if an “Interim Measures” pamphlet exists for your county and have one sent to you. You can also consult your local county extension office or pesticide state lead agency to determine whether they have imposed any requirements in your area to protect endangered and threatened species.

ENDANGERED SPECIES CONSIDERATIONS:

Whooping crane (*Grus americana*)

Territory: Wetlands; Migrant Eastern Montana

Least Tern (*Sterna antillarum*)

Territory: Yellowstone, Missouri River sandbars, beaches; Eastern Montana

Pallid Sturgeon (*Scaphirhynchus albus*)

Territory: Bottom dwelling; Missouri, Yellowstone Rivers.

White Sturgeon (Kootenai River population) (*Acipenser transmontanus*)

Territory: Bottom dwelling; Kootenai River

Ramik[®] Green should be placed in tamper-proof bait boxes or in areas which are inaccessible to children, pets, wildlife and domestic animals.

Wear rubber gloves when handling this product.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K₁, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin (Dicumerol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN

DIRECTIONS FOR USE

It is a violation of State and Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the package label for Ramik[®] Green (EPA Reg. No. 61282-46) must be in the possession of the user at the time of application. All applicable directions, restrictions, and precautions on the package label and on this supplemental label must be followed.

GROUND SQUIRRELS: (Columbian ground squirrel, *Spermophilus columbianus*) and (Richardson’s ground squirrel, *S. richardsoni*)

This product can be used to control ground squirrels found in levee and ditch banks, around farm buildings, along fence lines, in orchards, in crops, or in non-crop areas.

All outdoor bait placements must be made in protected bait stations, where they will be readily accessible to the ground squirrel, preferably near their burrows.

These bait stations must be strong enough to inhibit destruction by livestock and nontarget animals. Stations must be constructed to permit entry by target species but not by larger animals. Stations must be constructed to inhibit spillage of bait to the outside environment and must be secured so that they cannot be overturned by wind, livestock, pets and wildlife.

Bait placements should be checked frequently to insure that an uninterrupted supply of bait is always available. Maintain an uninterrupted supply of bait for at least 30 days or until all signs of feeding have stopped.

Bait stations may be maintained on a permanent basis to prevent reinfestation of the area.

Bait stations should be located conveniently to the squirrel burrows and should be placed 20 to 100 feet apart, depending on population density.

Use 2 to 4 pounds of bait in each protected bait station.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Plastic Containers: Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

Paper Containers: Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

NOTICE

Because Manufacturer and Seller have no control over storage, handling, and conditions of use, which are of critical importance, Manufacturer and Seller make no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate government agencies, due to misuse, improper handling or storage of this material. Nor do Manufacturer and Seller assume any responsibility for injury to persons, crops, animals, soil, or property arising out of misuse, improper handling, or storage of this material.

24(c) Registrant:

HACCO, Inc.

110 Hopkins Drive

Randolph, WI 53956