

CyKill™ Place Packs Kills Rats and Mice Breaks the Cycle

Norway Rats, Roof Rats, and House Mice usually consume a lethal dose in a single night's feeding, but it may take two or more days from the time of bait consumption for these rodents to die. Rodents cease feeding after consuming a toxic dose.

Kills anticoagulant resistant Norway Rats, Roof Rats & House Mice.

For Indoor and Outdoor (within 100 feet of buildings) control of Norway Rat, Roof Rat & House Mice

For Use in Attics, Kitchens, Basements, Garages, consistent with all use restrictions and label precautions and limitations.

{Text For Individual Container: It is illegal to sell placepacks individually.}

{Text For Container: Do not sell this product in individual containers holding less than 4 pounds of bait. }

ACTIVE INGREDIENT:	BY WT.
Bromethalin:	0.01%
OTHER INGREDIENTS:	99.99%
Total	100.00%

EPA Reg. No. 36488-52-61282

EPA Est. No. 61282-WI-01

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying the bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

FIRST AID

Have label with you when obtaining treatment advice.

If Swallowed	<ul style="list-style-type: none"> • Call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
---------------------	--

TREATMENT FOR PET POISONING

If an animal eats this bait, call a veterinarian at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the nerve poison Bromethalin. This is not an anticoagulant rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sub-lethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL! Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six (6) years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled or unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

USE RESTRICTIONS: This product can only be used to control Norway rats, roof rats and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Burrow baiting with this product is prohibited. Do not place near or inside ventilation duct opening. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA-inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than 4 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine where rats and mice will most likely find an opening and consume the bait. Generally, these areas are along walls, by gnawed openings or rub marks, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rodents have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Norway Rats and Roof Rats: Use 2 to 8 place packs per placement, using bait satiations as required under “**IMPORTANT**”. Make placements at intervals of approximately 15 to 30 feet at appropriate locations in infested area. Adjust the amount of bait provided according to the level of rat feeding expected at each placement location.

House Mice: Use 1 place pack per placement, using bait stations as required under “**IMPORTANT**”. Make placements at intervals of 8 to 12 feet at appropriate locations in infested areas. Up to 2 place packs may be placed at points of extremely high house mouse activity.

Baiting Strategy (Rats and Mice): Highest bait consumption is expected to occur on the first day or two after treatment begins. After several days, inspect the placement and replenish bait at sites where there is evidence of heavy feeding or where there is continued evidence of rodent activity. Normally, the volume of bait consumed will be significantly less with bromethalin baits than with anticoagulant baits. Although one night’s feeding will usually provide a lethal dose, maintain bait for at least one week to provide all rodents a chance to feed on the bait. Continue baiting until all signs of feeding have ceased.

Follow-Up: Replace contaminated or spoiled bait immediately. Using gloves, collect and properly dispose of all dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage and rodent-proof structures as much as possible. If reinfestation does occur, repeat treatment. When a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product, and spoiled and unconsumed bait, may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller’s liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For non-emergency (e.g. current product information)
call: **1-800-621-8829**

**Manufactured For:
HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956**

CyKill™ is a Trademark of Neogen Corporation

Optional Marketing Statements:

- Contains Bromethalin
- Palatable bait formulation
- Palatable formulation
- Ready to Use
- [Controls] [Kills] Norway rats, Roof rats, and House Mice
- [Controls][Kills] anticoagulant resistant Norway rats, Roof rats, and House Mice
- [Kills][Controls] Norway Rats, Roof Rats & House Mice in and around (within 50ft of) the home
- Kills][Controls] Norway Rats, Roof Rats & House Mice both inside and outside the home

Code (NI.RE.) NI.RE.

SPECIMEN LABEL

Item No.: Not Applicable
Net Contents: 0.75oz
UPC: Not Applicable
Format: UPC-A
Label: RS220441

Item No.: 112820
Net Contents: 100 x 0.75oz
UPC: 023626 30045 (3)
Format: UPC-A
Label: LA220825

Item No.: 112820
Net Contents: 4 x 100 x 0.75oz
UPC: 00023626 30065 (1)
Format: Interleaved 2-of-5
Label: Case Label