

Havoc[®] Rodenticide Bait Pack (Pellets)

Kills Rats and Mice

For Control of Commensal Rats and House Mice

Second Generation Anticoagulant

Kills Warfarin-Resistant Norway Rats and House Mice.

Norway Rats and house mice may consume a lethal dose in one night's feeding, with the first dead rodents appearing four or five days after feeding begins.

Distribution to and sales in "consumer" stores including grocery stores, drug stores, hardware stores, club stores is prohibited.

{Text For Individual Container: **INDIVIDUAL SALE PROHIBITED BY LAW**}

ACTIVE INGREDIENT:

Brodifacoum

3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one0.005%

OTHER INGREDIENTS¹:.....99.995%

TOTAL100.000%

¹ Contains Bitrex[®] brand denatonium benzoate

EPA Reg. No. 100-1056-61282

EPA Est. No. 61282-WI-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

HAVE LABEL WHEN OBTAINING TREATMENT ADVICE

If Swallowed	<ul style="list-style-type: none"> • Call a poison control center, doctor, or HOT LINE NUMBER immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin with plenty of water for 15-20 minutes. • Call a poison control center, doctor, or 1-800-498-5743 immediately for treatment advice.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal)

Call **1-800-498-5743**

For Chemical Emergency Assistance (Spill, Leak, Fire or Accident)

Call **1-800-424-9300**

NOTE TO PHYSICIAN: If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁ intramuscularly or orally. Vitamin K₁ is antidotal at doses of 10-20 mg (**not** mg/kg). Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or HOT LINE NUMBER at once.

NOTE TO VETERINARIAN

Anticoagulant Brodifacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K₁ at doses of 2-5 mg/kg. if needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin or on clothing. All handlers (including applicators must wear long-sleeve shirt, long pants, shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may also be hazardous to aquatic organisms in water adjacent to treated areas, do not contaminate water when disposing of equipment wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To reduce exposure:

- Store product not in use in a location out of reach of children and pets. When used in residential structures this product must be applied in tamper resistant bait stations. All above ground, outdoors applications must be in tamper resistant bait stations. These stations must be resistant to destruction by dogs and by children under 6 years of age, and must be used in a manner that prevents such children from

reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.

- Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice in and around homes, industrial, commercial, agricultural and public buildings, and similar manmade structures. **Do not apply further than 50 feet from buildings.** Do not place inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait.

Do not sell this product in individual containers holding less than 8 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

Norway and Roof Rats:

Apply 2-8 place packs (usually at intervals of 15-30 ft.) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. Two place packs contain enough bait (100 g) to kill 1-3 rats.

House Mice:

Open pack and apply ¼ - ½ oz. of bait per placement. Space placements at intervals of 8-12 ft. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Follow-Up: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and (when baiting operations are to be ceased) unconsumed bait according to the **Disposal** paragraph. To prevent reinfestation, limit food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish or maintain bait stations and replenish with census or toxic bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Dispose of wastes resulting from the use of this product at an approved waste disposal facility or call your local solid waste agency for alternative disposal instructions. Never dispose of unused product down any indoor or outdoor drain.

Container Handling (pail; pouch; and outer box): Nonrefillable container. Do not reuse or refill this container. When completely empty, offer for recycling if available, or dispose of empty container in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

Hacco, Inc. and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Hacco, Inc., and Buyer and User assume the risk of any such use. **HACCO, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

In no event shall Hacco, Inc. or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF HACCO, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE HACCO, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SPECIMEN LABEL

Hacco, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Hacco, Inc.

For non-emergency (e.g., current product information) call:
1-800-621-8829

Manufactured For:
HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956

Havoc[®] is a trademark of a Syngenta Group Company.
Bitrex[®] is a registered trademark of Macfarlan Smith Ltd.,
Edinburgh, Scotland.
Product of the United Kingdom
Formulated in the USA

Code (NG.RN.) NR.RR.

Item No.: 116372
Net Contents: 40 x 2 x 50 g
UPC: 095242 26372 (6)
Format: UPC-A
Label:

Item No.: 116374
Net Contents: 240 x 2 x 50 g (52.8 lbs)
UPC: 00095242 16374 (3)
Format: Interleaved 2-of-5
Label:

Item No.: 116381
Net Contents: 12 x 50 g (1.76 oz.)
UPC: 095242 16375 (0)
Format: UPC-A
Label: